

# EVIDENCE FOR A 13,14-CIS CYCLE IN BACTERIORHODOPSIN

PAUL TAVAN AND KLAUS SCHULTEN

*Physik-Department, Technische Universität München, 8046 Garching, Federal Republic of Germany*

**ABSTRACT** We discuss to what extent the vibrational spectra of bacteriorhodopsin that have been observed and assigned by Smith et al. (1, 2) by means of resonance Raman and by Gerwert and Siebert (*EMBO (Eur. Mol. Biol. Organ.) J. In press*) by means of infrared absorption experiments are in agreement with a photo-cycle of bacteriorhodopsin that involves the sequence BR, IO(all-*trans*)  $\rightarrow$  K(13,14-*cis*)  $\rightarrow$  L(13,14-*cis*)  $\rightarrow$  M(13-*cis*)  $\rightarrow$  N(13-*cis*)  $\rightarrow$  O(all-*trans*). Our discussion is based on a quantumchemical modified neglect of diatomic overlap [MNDO] calculation of the vibrational spectra of the relevant isomers of the protonated retinal Schiff base. In particular, we investigated in these calculations the effects of different charge environments on the frequencies of the relevant C-C single bond stretching vibrations of these isomers.

## INTRODUCTION

Bacteriorhodopsin (BR) in the cell membrane of *Halobacterium halobium* works as a light-driven proton pump (3-5). In 1978 we proposed a molecular mechanism for the photo-cycle of BR which was based on a primary all-*trans*  $\rightarrow$  13,14-*cis* photoisomerization reaction of the chromophore (6). In our subsequent work we carried out quantumchemical calculations on the chromophore, i.e., protonated and unprotonated retinal Schiff bases in various isomeric states and in different charge environments. These calculations provided strong arguments for our proposition (7-10). Recently, the suggested mechanism has been challenged by Smith et al. (1) on the basis of calculations that relate geometries and observed vibrational spectra of polyenes. The authors used an empirical force field in their calculations (11). An alternative approach is to employ for the vibrational analysis a force field evaluated from quantum chemical calculations. In this article we will follow this approach. We will show that the resonance Raman (RR) data of Smith et al. (2, 12) and other vibrational spectra (Gerwert, K., and F. Siebert, *EMBO (Eur. Mol. Biol. Organ.) J. In press.*) can be reconciled with our suggestion of a 13,14-*cis* intermediate in the BR pump cycle.

The arguments that lead us to postulate a 13,14-*cis* cycle in BR had been based on the observations that the initial state BR<sub>568</sub> contains all-*trans* retinal (13-16) as a protonated Schiff base which becomes deprotonated and is 13-*cis* in the long-lived M<sub>412</sub> intermediate (14-17). The arguments are the following (6, 18, 7, 8): (a) A primary process all-*trans*  $\rightarrow$  13,14-*cis* involves up to the transition state the motion of only small masses and, therefore, can be very fast. This argument is corroborated by the recent finding of Nuss et al. (19) that the primary process in BR is

completed after  $\sim$ 430 fs. (b) A 13,14-*cis* intermediate is thermally stable against the direct back-isomerization, whereas a 13-*cis* intermediate is not (9). The fact that thermal reversals of the primary reaction have not been observed in BR and would, indeed, be detrimental to the pump efficiency, supports our suggestion. (c) An indirect, i.e., sequential reisomerization 13,14-*cis*  $\rightarrow$  13-*cis*  $\rightarrow$  all-*trans* enforces a deprotonation and a subsequent reprotonation of the chromophore. Such deprotonation and reprotonation, in fact, are observed in the cycle.

The arguments of Smith et al. (1) against our mechanism are derived from their observations of the RR spectra of the intermediates in the bacteriorhodopsin cycle. The authors substituted the natural carbons <sup>12</sup>C at the C<sub>14</sub> and the C<sub>15</sub> positions of retinal by <sup>13</sup>C and determined which of the vibrational bands are affected by this substitution. In this way they could assign the C<sub>14</sub>-C<sub>15</sub> stretch vibration. This vibration is very sensitive to geometrical changes of the chromophore and, hence, the assignment of this band for the intermediates of the BR cycle implies information on the isomeric state of retinal. However, the only key to this information are calculations of vibrational spectra.

Smith et al. determined that the C<sub>14</sub>-C<sub>15</sub> stretch frequency shifts from 1,201 cm<sup>-1</sup> for Br<sub>568</sub> to 1,195 cm<sup>-1</sup> for the low temperature (77°K) primary BR photoproduct K<sub>625</sub> (1) and 1,172 cm<sup>-1</sup> for the L<sub>550</sub> intermediate (2). Gerwert and Siebert, on the basis of infrared difference spectroscopy (IR) using the same isotopic labels as Smith et al. (2) corroborated the K<sub>625</sub> assignment (20) but assigned the C<sub>14</sub>-C<sub>15</sub> stretch frequency for L<sub>550</sub> to a band at 1,155 cm<sup>-1</sup> (*EMBO (Eur. Mol. Biol. Organ.) J. In press.*). To extract a statement on retinal's geometry from these vibrational shifts, Smith et al. (1) quoted a calculation by Curry (21) on butadiene, which had indicated that the stretch frequency of the central single bond of this com-

pound experiences a downshift by  $\sim 50$  to  $100\text{ cm}^{-1}$  upon rotation around this single bond. Furthermore, these authors carried out a similar calculation on a protonated Schiff base fragment of the size of butadiene rendering a comparably large red shift ( $69\text{ cm}^{-1}$ ) of the single bond stretch frequency upon *s-trans*  $\rightarrow$  *s-cis* isomerization. All these calculations required to fit a large number of parameters for an intramolecular force field (22), in order to describe with a high degree of accuracy vibrational spectra of protonated retinal Schiff bases (RSBH<sup>+</sup>) (11). Since (a) the result on the Schiff base fragment matched the butadiene result rather well; (b) the C<sub>13</sub>-C<sub>14</sub>-C<sub>15</sub>-N portion of the all-*trans* RSBH<sup>+</sup> corresponds geometrically to *s-trans* butadiene, the corresponding 13,14-*cis* compound to *s-cis* butadiene; and (c) the C<sub>14</sub>-C<sub>15</sub> stretch frequency of retinal does not experience a large downshift upon the Br<sub>568</sub> to K<sub>625</sub> transformation, the authors determined that a 13,14-*cis* geometry must be excluded for the K intermediate. The argument hinges on the validity of intramolecular forces obtained by fitting indiscriminately the vibrational spectra of polyenes and retinal Schiff bases with different  $\lambda_{\text{max}}$  values, i.e., hinges on the similarity of force constants between butadiene and the C<sub>13</sub>-C<sub>14</sub>-C<sub>15</sub>-N segment of RSBH<sup>+</sup> in the far red-shifted intermediate K<sub>625</sub>. We will show below that such similarity does not exist and that the force fields as witnessed by vibrational spectra can vary sensitively with the  $\lambda_{\text{max}}$  values of retinal compounds.

## METHODS

To provide a parameter-free, nonempirical description of the vibrational modes of the BR chromophores we have carried out quantumchemical MNDO force calculations for the vibrational spectra of protonated model compounds representing the complete polyene part of the RSBH<sup>+</sup>.

We have chosen the MNDO method (modified neglect of diatomic overlap) (23) as it furnishes a quantitatively correct description of the ground state properties of protonated retinal Schiff bases. This has been demonstrated by us in our quantumchemical investigation of the effects of protonation and electrical interactions (9) and of substituents (10) on the stereochemistry of retinal Schiff bases. That MNDO gives a correct account of the changes induced by protonation in the bonding properties of molecules containing large  $\pi$ -systems has been demonstrated by us also for the merocyanine dye stilbazolium betaine (24), in which a thermal isomerization around the central single  $\pi$ -bond is prevented in highly polar, protic solvents (25).

For the vibrational analysis we used the MOPAC program package distributed by the Quantum Chemistry Program Exchange of Indiana University at Bloomington (QCPE 455). In a typical calculation of the vibrational spectrum for a model RSBH<sup>+</sup> like one of those depicted in Fig. 1 A-E, first the ground state equilibrium geometry is calculated by minimizing the MNDO total molecular energy  $E(Q_i)$  with respect to essentially all nuclear coordinates  $Q_i$ ,  $i = 1, \dots, 3N-6$ , where  $N$  is the number of atoms in the molecule. Next, the full Hessian matrix of second derivatives  $\partial^2 E / \partial Q_i \partial Q_k$  with respect to the Cartesian coordinates  $Q_i$  is evaluated. Multiplication of this MNDO force matrix with the mass weights and diagonalization yields then the vibrational modes in Cartesian coordinates and the corresponding vibrational frequencies. The assignment of these modes to local internal coordinates like "C<sub>14</sub>-C<sub>15</sub> stretch" or "C<sub>14</sub>-H rock," etc., has been achieved by a graphical representation and visual inspection. To provide some examples, Fig. 1 A-E show the C<sub>14</sub>-C<sub>15</sub> stretching mode of the model RSBH<sup>+</sup> for various chromophore geometries. The assignment has been checked by the

investigation of suitable isotope effects and by the creation of motion pictures on a high resolution graphics screen.

As has been observed previously for the complete neglect of diatomic overlap (CNDO/2) method (26) also, the MNDO force field overestimates systematically all force constants except (see below) those for the C-C single bond torsions. But as in the case of CNDO/2 this overestimate can be corrected by an appropriate scaling of the force matrix, yielding then a satisfactory description of normal modes and vibrational frequencies. This has been demonstrated by us for the well-known vibrational spectra of butadiene and hexatriene and their deuterated derivatives (Tavan, P., Z. Luo, and K. Schulten, unpublished results). The necessary scaling of the MNDO force matrix is accomplished in our method by the introduction of MNDO "spectroscopic" masses. The choice of 1.094 for the hydrogen mass and of 14.75 for the carbon atoms turned out to reproduce the observed butadiene spectrum, for instance, with absolute errors of at most  $30\text{ cm}^{-1}$  in a spectral range between 0 and  $2,000\text{ cm}^{-1}$ . The only exception is the torsional vibration around the central single bond which is calculated at  $41\text{ cm}^{-1}$  and observed at  $163\text{ cm}^{-1}$  (26). This failure rests on the qualitatively incorrect description of the torsional stability of long C-C bonds by MNDO and related quantumchemical methods which we have discussed in detail in reference 9. Due to this error the dihedral angles of C-C single bonds in unpolar compounds like the protonated retinal Schiff bases with a close ( $2\text{ \AA}$ ) counter ion have to be excluded from the MNDO geometry optimization (9). However, the important spectral features of polyene dyes and their derivatives are rather well covered by NDO force calculations like MNDO or CNDO/2, since the low frequency single bond torsional modes couple only weakly to the high frequency hydrogen out of plane wags, C-C single and double bond stretches and hydrogen in plane rocks, (cf. reference 26 and references quoted therein). Hence, we have chosen planar C-C single bond geometries for all isomers of the model RSBH<sup>+</sup> with a close counter ion (cf. Fig. 1 A, B, and E) except for the 13,14-*cis* isomer, in which a nonplanar geometry is enforced by intramolecular sterical hindrance. To determine a reasonable model geometry for this isomer, we have calculated first the equilibrium geometry for the highly polar 13,14-*cis* cation relaxing all degrees of freedom, since in such a compound the above-mentioned error is absent (9). MNDO predicted the cation to be twisted around the C<sub>13</sub>-C<sub>14</sub> bond by  $16^\circ$  and around the C<sub>14</sub>-C<sub>15</sub> bond by  $22^\circ$  (cf. Fig. 1 D). As the torsional stability of the C<sub>13</sub>-C<sub>14</sub> double bond increases in the less polar compounds, we have chosen a torsional angle of  $45^\circ$  for the C<sub>14</sub>-C<sub>15</sub> bond in the 13,14-*cis* isomers with a counter ion (cf. Fig. 1 C). It turned out that the calculated vibrational spectra are insensitive to small ( $\approx 10^\circ$ ) changes of these dihedral angles.

As we have shown in reference 9, the model retinal Schiff bases depicted in Fig. 1 represent the electronic properties of the complete chromophore very well. They should also provide a good description for those vibrational modes that are localized in the polyene part of the molecule toward the Schiff base nitrogen. For most isomeric states and charge environments the C<sub>10</sub>-C<sub>11</sub> and C<sub>14</sub>-C<sub>15</sub> single bond stretches turned out to be rather well localized in this region. To exclude an artificial coupling of the in-plane bending modes of the two hydrogen atoms at C<sub>5</sub> and C<sub>6</sub> of the model compound we have decoupled the motion of these atoms by assigning them arbitrarily a large mass of 15. The effect of this decoupling amounted in a nearly uniform shift of the single bond stretches of  $\sim 5\text{ cm}^{-1}$ . For the same reason the counter ions have been spatially fixed, assigning them a very large mass of 1,000.

## RESULTS

We have applied the calculations described in Methods to determine the C<sub>10</sub>-C<sub>11</sub> and the C<sub>14</sub>-C<sub>15</sub> stretch frequencies of the RSBH<sup>+</sup>. These vibrations were selected as they constitute prominent bands in the RR and IR difference spectra of BR. The question that we investigated first was to what extent these vibrations are sensitive to the isomeric




FIGURE 1 The MNDO equilibrium geometry and the  $C_{14}$ - $C_{15}$  stretch normal mode of the model RSBH<sup>+</sup> in the field of a negative charge and of a  $Cl^-$  ion 2 Å distant from the proton at the Schiff base nitrogen is shown for four different isomeric states: (A) all-*trans*, (B) 13-*cis*, (C) 13,14-*cis*, (E) 13,15-*cis*. The negative charge 3 Å distant from  $C_5$  and perpendicular to the plane of the molecule has been omitted in the drawing. Also shown are the geometry and the  $C_{14}$ - $C_{15}$  stretching mode for the 13,14-*cis* undisturbed cation (D).

state, to the protonation state, and to interactions with external charges.

Our results are summarized in Table I. This table provides the vibrational frequencies for the relevant geometries of retinal: all-*trans*, 13-*cis*, 13,14-*cis*, 13,15-*cis*. The model retinal Schiff base was chosen protonated with a  $\text{Cl}^-$  ion as counter ion at a distance of 2 Å and 3 Å near the proton at the terminal nitrogen and with no counter ion. In addition, a negative charge has been located at a distance of 3 Å near  $\text{C}_5$ . The first compound represents a chromophore with a spectrum in the blue, i.e., at ~480 nm, the second compound a chromophore with a spectrum at ~550 nm, and the last compound a chromophore with a spectrum at 640 nm. To demonstrate the influence of the negative charge near  $\text{C}_5$  we have also included the data on the pure cations representing chromophores with spectra at ~600 nm (27, 28). Therefore, in comparisons with experimental data, we will take the 2-Å compounds as models for protonated Schiff bases in solution, the 3-Å compounds as

TABLE I  
THE MNDO-FORCE  $\text{C}_{14}\text{--C}_{15}$  AND  $\text{C}_{10}\text{--C}_{11}$  STRETCH FREQUENCIES FOR THE MODEL  $\text{RSBH}^+$

Compound	$\text{C}_{14}\text{--C}_{15}$			$\text{C}_{10}\text{--C}_{11}$		
	$^{12}\text{C}, ^1\text{H}$	$^{13}\text{C}$	$^2\text{H}$	$^{12}\text{C}, ^1\text{H}$	$^{13}\text{C}$	$^2\text{H}$
	$\text{cm}^{-1}$	$\text{cm}^{-1}$	$\text{cm}^{-1}$	$\text{cm}^{-1}$	$\text{cm}^{-1}$	$\text{cm}^{-1}$
all- <i>trans</i>						
2 Å, C	1,222.8	1,207.4	1,224.1	1,178.6	1,177.4	1,183.3
3 Å, C	1,229.6	1,209.7	1,229.4	1,179.3	1,177.3	1,187.3
cat., C	1,268.9	1,259.9		1,207.6	1,205.1	
cat.	1,215.4	1,207.2		1,182.3	1,180.3	
13- <i>cis</i>						
unprot.	1,218.6			1,188.1		
2 Å, C	1,208.7	1,201.3	1,216.3	1,189.4	1,183.6	1,189.6
3 Å, C	1,212.8	1,203.8		1,194.2	1,188.5	
cat., C	1,236.2	1,228.5		1,220.9	1,219.0	
cat.	1,226.5	1,214.8		1,204.2	1,201.1	
13- <i>cis</i> , 14s- <i>cis</i>						
2 Å, C	1,138.8	1,125.0		1,196.4	1,194.7	
3 Å, C	1,168.1	1,143.2		1,202.9	1,199.3	
cat., C	1,230.1	1,227.3		1,217.8	1,214.1	
cat.	1,218.0	1,209.0		1,185.0	1,183.2	
13,15- <i>cis</i>						
2 Å, C	1,155.1		1,211.5	1,193.9		1,190.6
3 Å, C	1,182.4		1,229.9	1,195.7		1,195.5
cat., C	1,215.6	1,204.5		1,220.7	1,219.9	
cat.	1,205.4		1,240.0	1,189.7		1,190.6

The MNDO-force  $\text{C}_{14}\text{--C}_{15}$  and  $\text{C}_{10}\text{--C}_{11}$  stretch frequencies for the model  $\text{RSBH}^+$  are given for various isomeric states, charge environments, and isotopic modifications. The column labels specify the isotopic substitutions considered. The column labeled by  $^{12}\text{C}, ^1\text{H}$  gives the frequencies for the unsubstituted compounds. The  $^{13}\text{C}$  column gives those for the 14,15- $^{13}\text{C}$  isotopes. The  $^2\text{H}$  column gives those for the deuterated Schiff bases. The charge environment is characterized by the letter "C" if a negative charge was placed 3 Å distant to atom  $\text{C}_5$ . It is characterized by the distances "2 Å" or "3 Å" if a  $\text{Cl}^-$  ion was positioned at that distance from the proton at the nitrogen, and by the keyword "cat." if no counter ion was present in the calculation.

models for BR chromophores absorbing ~550 nm, and the cations with and without negative charge near  $\text{C}_5$  as models for the  $\text{K}_{625}$  and  $\text{O}_{640}$  intermediates. The above choice of model charge environments had been motivated by the experimental investigation of the BR binding site by Derguini et al. (29), by the corresponding theoretical analysis in terms of an external point charge model (30, 31), and by our own analysis of the thermal isomerization properties of retinal in BR (9).

We have identified the vibrational modes in Table I by visual inspection of all the eigenmodes of the chromophores. Those stretching vibrations most localized around the 10–11 and 14–15 bonds were included. However, to prove that the vibration selected are indeed identical to the ones identified in the experimental spectra we applied the same criteria as the experimentalists and for this purpose determined how the frequencies are shifted upon isotopic substitutions, i.e., either  $^{13}\text{C}$  substitution at the  $\text{C}_{14}$  and the  $\text{C}_{15}$  positions or substitution of N–H by N–D. The reader may consult the corresponding entries of Table I when we make comparison between our calculated vibrational frequencies and those observed by Smith et al. (2, 12) and by Gerwert and Siebert (*EMBO (Eur. Mol. Biol. Organ.) J. In press.*).

### Structure of $\text{Br}_{548}$

When our earlier quantumchemical calculations revealed that an  $\text{RSBH}^+$  can thermally undergo concerted isomerizations around two double bonds, i.e., all-*trans* → 13,15-*cis*, we suggested (7) that the dark-adapted pigment  $\text{Br}_{548}$  may exist in a 13,15-*cis* geometry. The observation by Oesterhelt et al. (13) that an all-*trans* → 13-*cis* isomerization takes place during dark-adaptation had been the experimental background for this suggestion. Fig. 1 *A* and *E*, shows that the geometry of 13,15-*cis* retinal is very close to that of the all-*trans* compound, and that one may consequently expect that both compounds are about equally stable in the BR binding site.

The suggestion of a 13,15-*cis*  $\text{Br}_{548}$  compound was proven to be correct by Smith et al. (2) on the basis of RR and by Harbison et al. (32) on the basis of nuclear magnetic resonance (NMR) spectra. Here we want to show that an application of our own results to the RR spectra of  $\text{Br}_{568}$  and  $\text{Br}_{548}$  reproduces this assignment. For this purpose we present in Fig. 2 the relevant data taken from Table I as well as the experimental observations (2). These data represent the effects of deuteration of the Schiff base nitrogen on the  $\text{C}_{14}\text{--C}_{15}$  stretch frequency. The calculations show that a C–N bond with a *trans* geometry of the nitrogen proton relative to the hydrogen at  $\text{C}_{14}$  (see Fig. 1 *A*) does not exhibit a frequency shift of the  $\text{C}_{14}\text{--C}_{15}$  stretch upon deuteration. However, due to the coupling of the N–H rocking mode to the  $\text{C}_{14}\text{--C}_{15}$  stretch, a C–N bond with a proton in the *cis*-position (see Fig. 1 *E*) reveals a rather strong deuteration effect, i.e., an upshift of 48  $\text{cm}^{-1}$ . Fig. 2 shows that the  $\text{C}_{14}\text{--C}_{15}$  stretch frequency of  $\text{Br}_{568}$




FIGURE 2 The effect of deuteration of the Schiff base nitrogen on the  $C_{14}$ - $C_{15}$  stretch frequency calculated for the model RSBH<sup>+</sup> in the field of a  $Cl^-$  ion at a distance of 3 Å is compared with the experimental findings by Smith et al. (2).

does not exhibit a deuteration effect, whereas the  $C_{14}$ - $C_{15}$  stretch frequency of  $Br_{548}$  does; its shift upon deuteration measures  $41\text{ cm}^{-1}$ . Such an analogy between calculation and observation led Smith et al. (2) to assign the 13,15-*cis* geometry (with a *cis*-proton at the C-N bond) to  $Br_{548}$ . These authors calculated an upshift of 5 to  $12\text{ cm}^{-1}$  for a *trans* C-N bond and an upshift of  $62\text{ cm}^{-1}$  for a *cis* C-N bond. Fig. 2 demonstrates that our calculations are in harmony with the results of Smith et al. (2), reproducing the behavior of the pigments  $Br_{568}$  and  $Br_{548}$  even somewhat better.

### Effect of Electrical Interactions on the Vibrational Spectra of the Isomers of RSBH<sup>+</sup>

In this section we want to demonstrate that, in addition to the geometry, electrical interactions also have an important effect on the vibrational spectra of some of the retinal isomers. Before presenting our results we wish to point out that electrical interactions and their variation during the pump cycle play an important role in BR, a role which can be easily witnessed from the strong spectral shifts of the pigment (28).

In Fig. 3 A we present the  $C_{14}$ - $C_{15}$  stretch frequency for the model RSBH<sup>+</sup> in the all-*trans*, the 13-*cis*, the 13,15-*cis*, and the 13,14-*cis* isomeric state. The electrical interactions considered correspond to an RSBH<sup>+</sup> without a counter ion and with a counter ion at distances of 2 Å and 3 Å. All these compounds are subjected to the field of a negative charge near  $C_5$ . The effect of removal of this charge is represented by the data on the pure RSBH<sup>+</sup> cation. The results in Fig. 3 A are the main results of this paper.

Fig. 3 A demonstrates that an alteration of the electrical interactions has a considerable effect on all the isomers.




FIGURE 3 The effect of a varying charge environment on (A) the  $C_{14}$ - $C_{15}$  and (B) the  $C_{10}$ - $C_{11}$  stretch frequency is shown for four isomers of the model RSBH<sup>+</sup>. The charge environment is indicated by the distance of a possible  $Cl^-$  ion to the Schiff base proton and by the letter C indicating the presence of a negative charge 3 Å near  $C_5$ . Related chromophore structures are connected by dotted lines. Also given are the observed values for  $Br_{568}$  (12),  $Br_{548}$  (2),  $L_{550}$  (Gerwert, K., and F. Siebert, *EMBO (Eur. Mol. Biol. Organ.) J.* In press.), for an all-*trans* and 13-*cis* RSBH<sup>+</sup> in solution (33) and our tentative assignment of bands in the spectra of Alshut and Stockburger (personal communication) to the N intermediate (see Results, Mechanism of the Proton Pump for discussion).

This effect is smallest in the case of the 13-*cis* compound. The effect of electrical interactions is only a little stronger for the all-*trans* compound. However, in the case of the 13,15-*cis* compound, the different charge environments induce large shifts. These shifts are found to be even larger for the 13,14-*cis* compound.

To rationalize the strong effects of external charges on the  $C_{14}$ - $C_{15}$  stretch frequency in the isomers, we present in Table II a comparison of the bond lengths of the four 13,14-*cis* protonated retinal Schiff bases considered so far: the compounds in the field of the negative charge near  $C_5$ , with a  $Cl^-$  counter ion at a distance of 2 Å and 3 Å, or without a  $Cl^-$  and the pure cation, i.e., without the charge and without a  $Cl^-$ . Table II shows that the single and

TABLE II  
THE MNDO EQUILIBRIUM BOND LENGTHS OF THE  
13,14-*cis* MODEL RSBH<sup>+</sup>

Bond description	2 Å, C	3 Å, C	cat., C	cat.
	Å	Å	Å	Å
Single bonds				
C <sub>6</sub> —C <sub>7</sub>	1.463	1.462	1.448	1.461
C <sub>8</sub> —C <sub>9</sub>	1.477	1.474	1.449	1.469
C <sub>10</sub> —C <sub>11</sub>	1.460	1.455	1.418	1.440
C <sub>12</sub> —C <sub>13</sub>	1.467	1.460	1.415	1.437
C <sub>14</sub> —C <sub>15</sub>	1.461	1.452	1.396	1.414
Double bonds				
C <sub>5</sub> —C <sub>6</sub>	1.355	1.355	1.364	1.354
C <sub>7</sub> —C <sub>8</sub>	1.358	1.360	1.377	1.364
C <sub>9</sub> —C <sub>10</sub>	1.370	1.373	1.402	1.383
C <sub>11</sub> —C <sub>12</sub>	1.363	1.368	1.405	1.386
C <sub>13</sub> —C <sub>14</sub>	1.370	1.377	1.424	1.405
C—N bond				
N <sub>15</sub> —C <sub>16</sub>	1.315	1.320	1.359	1.347

The MNDO equilibrium bond lengths of the 13,14-*cis* model RSBH<sup>+</sup> are given as a function of the charge environment (see the caption to Table I for an explanation).

double bonds near the C—N bond are most strongly affected by a change in the charge environment as represented by these four compounds. For example, the length of the C<sub>14</sub>—C<sub>15</sub> single bond varies between 1.46 and 1.40 Å. The effect on the bond length develops mainly upon removal of the counter ion, i.e., corresponds to the difference between the 3-Å compound and the compounds without counter ion. The negative charge near C<sub>5</sub> amplifies the effect of protonation, as is revealed by a comparison of the geometries of the two cations: the cation in the field of the charge exhibits nearly equal single and double bond lengths in the part of the carbon chain between atoms C<sub>9</sub> and C<sub>15</sub>. This compound resembles more a cyanine dye than a polyene with bond alternation. In contrast, the pure cation still shows some degree of bond alteration in the region between C<sub>9</sub> and C<sub>13</sub>.

One expects that the alteration of a bond length reflects a change in the force constant of this bond and, therefore, that the variation of the bond length should be accompanied by an alteration in the stretch frequency. This is, in fact, the behavior shown in Fig. 3A. However, for a complete understanding of the calculated vibrational frequencies, one has to also consider the coupling of the bond stretch motions to other degrees of freedom. This coupling as well as the frequency distribution of the other degrees of freedom is also subject to change upon alterations of the charge environment. Furthermore, this coupling may differ in the various isomeric states. Hence, the effect of a charge redistribution on the stretch frequencies cannot be attributed to a single cause (force constant), but rather is the result of a complex interplay of several causes (force constants, kinetic couplings).

Fig. 3 reveals also how the C<sub>14</sub>—C<sub>15</sub> stretch frequency

changes solely due to geometry changes when the charge environment is not altered. As an example we may consider the nonpolar 2-Å compound. Its C<sub>14</sub>—C<sub>15</sub> stretch frequency downshifts by only 14 cm<sup>-1</sup> upon an all-*trans* → 13-*cis* isomerization. For retinal Schiff bases in solvents one observes a value of 15 cm<sup>-1</sup> (1,11,33). An all-*trans* → 13,15-*cis* isomerization has a stronger effect, downshifting the frequency by 45 cm<sup>-1</sup>. The largest effect is found for the all-*trans* → 13,14-*cis* isomerization, which lowers the C<sub>14</sub>—C<sub>15</sub> stretch frequency by 84 cm<sup>-1</sup>. This latter shift is in agreement with the aforementioned behavior of butadiene when it undergoes isomerization around its central single bond. In the butadiene case, MNDO predicts a downshift of 99 cm<sup>-1</sup>.

A most remarkable feature of the frequency behavior shown in Fig. 3A is the invariance of the C<sub>14</sub>—C<sub>15</sub> stretch frequency of the compounds without counter ion (cations). The C<sub>14</sub>—C<sub>15</sub> stretch frequency in these compounds exhibits a behavior that is completely different from its behavior in the nonpolar compounds. None of the isomerization processes affects the C<sub>14</sub>—C<sub>15</sub> stretch frequency in a significant way. The reasons for this behavior are opposing effects from geometry changes and changes in the electronic structure. The difference in the behavior of the cation compounds and the 2-Å compound is manifested by the fact that the C<sub>14</sub>—C<sub>15</sub> stretch frequency of the 13,14-*cis* pure cation lies *above* that of the all-*trans* pure cation, whereas this frequency in the 13,14-*cis* 2-Å compound experiences a very strong downshift compared with its all-*trans* relative. Furthermore, the C<sub>14</sub>—C<sub>15</sub> stretch frequencies of both 13,14-*cis* cations lie only ~8 cm<sup>-1</sup> below those of the corresponding 13-*cis* compounds, whereas very large downshifts of 70 and 45 cm<sup>-1</sup> upon 14*s-trans* → *s-cis* isomerization are predicted for the more nonpolar, polyene-like 2 Å and 3-Å 13-*cis* compounds, respectively. This dramatic difference, which is not predicted by the empirical calculations of Smith et al., provides our key argument against the geometric assignment of these authors (reference 1): the empirical force field employed by these authors may be of some use for the description of nonpolar molecules as it has been derived from small polyene-like compounds. But it fails in the description of the highly polar, far red-shifted chromophores.

Fig. 3B shows that the C<sub>10</sub>—C<sub>11</sub> stretch frequency exhibits only a minor dependence on charge alteration and isomerizations around the three terminal bonds.

#### Assignment of a 13,14-*cis* Intermediate

To use the results of Fig. 3A we note that the primary intermediate K of the pump cycle in BR has a strongly red-shifted absorption spectrum and should, therefore, correspond to one of the cation compounds. This correspondence implies, then, according to Fig. 3A, that the C<sub>14</sub>—C<sub>15</sub> stretch frequency should not allow a differentiation between a 13-*cis* and a 13,14-*cis* geometry for the K intermediate. This is in disagreement with Smith et al. (1),

who claimed that they can assign a particular one of these geometries to this intermediate by comparing the very small downshift of  $6\text{ cm}^{-1}$  observed in the transition from  $\text{BR}_{658}$  to  $\text{K}_{625}$  with the frequency shift of butadiene: the authors claimed that a  $50$  to  $100\text{ cm}^{-1}$  downshift should have been observed relative to all-*trans*  $\text{Br}_{568}$  if the K intermediate were to assume a 13,14-*cis* geometry.

Fig. 3 A suggests, however, that one should look at the L intermediate for a decision between a 13-*cis* and a 13,14-*cis* geometry. Such decision can be derived since  $\text{Br}_{568}$ ,  $\text{Br}_{548}$ , and  $\text{L}_{550}$  have all similar spectra which should correspond to the 3-Å compounds in Fig. 3 A. The fact that the  $\text{C}_{14}\text{--}\text{C}_{15}$  stretch frequency of the L intermediate lies  $46\text{ cm}^{-1}$  below the corresponding frequency of  $\text{Br}_{568}$  (all-*trans*) and still  $12\text{ cm}^{-1}$  below that of  $\text{Br}_{548}$  (13,15-*cis*) should imply that the L intermediate is 13,14-*cis*. This argument is strongly supported by the assignment of Gerwert and Siebert (*EMBO (Eur. Mol. Biol. Organ.) J. In press.*) of the  $\text{C}_{14}\text{--}\text{C}_{15}$  stretch frequency at  $1,155\text{ cm}^{-1}$  for L. The suggestion of Smith et al. of a 13-*cis* geometry for L cannot be reconciled with the results of Fig. 3 A. Even the previous assignment of a band at  $1,172\text{ cm}^{-1}$  to the  $\text{C}_{14}\text{--}\text{C}_{15}$  stretch by Smith et al. (2) appears to be at variance with a 13-*cis* conformation of L because a 13-*cis* RSBH<sup>+</sup> in solution has a  $\text{C}_{14}\text{--}\text{C}_{15}$  stretch frequency of  $1,175\text{ cm}^{-1}$  (33) and because one has to expect an upshift by  $\sim 10\text{ cm}^{-1}$  upon incorporation into the protein (11). In our calculations this "opsin" shift is reflected in the general upshift of the stretch frequencies of the 3-Å compounds as compared with the 2-Å compounds. Assuming that no isomerization takes place in the  $\text{K} \rightarrow \text{L}$  transition we assign also a 13,14-*cis* geometry to K.

### Mechanism of the Proton Pump

In this final section we would like to summarize our results in view of a mechanism of the pump cycle in BR. The pump cycle is presented in Fig. 4. We will first provide the arguments that lead to this pump cycle and demonstrate then by means of Fig. 5 that the calculated  $\text{C}_{14}\text{--}\text{C}_{15}$  stretch frequencies of the isomers occurring in the cycle are in harmony with the available data.

The cycle in Fig. 4 presents the terminal part of the retinal chromophore and three acids  $\text{A}_i$  which alter their protonation state during the cycle. The protonation states of the three groups assumed here were suggested by Engelhard et al. (34) on the basis of their difference IR spectra. These authors also claim that the three groups  $\text{A}_i$  shown are aspartic acids. The protonation changes of these acids are connected with either the approach or the release of protons from the vicinity of retinal, as indicated in Fig. 4. If protons released and absorbed are originating from different sides of the purple membrane, the reaction sequence of Fig. 4 can obviously explain a vectorial proton pump.

The geometries of retinal assumed in the cycle of Fig. 4 are those that we suggested since our first proposal (6).




FIGURE 4 The 13,14-*cis* model for the BR proton pump cycle. For each intermediate the chromophore conformation and the Schiff base charge environment are indicated schematically. Note that the pump cycle as described transports two protons when O is involved and one proton when O is bypassed.

What is new in the representation of Fig. 4 are the relative positions of the acids  $\text{A}_i$  and retinal. The positions have been chosen to rationalize the spectra of retinal observed at the different stages of the cycle (28). In the  $\text{Br}_{568}$  stage the Schiff base nitrogen forms a salt bridge, i.e., sees a close counter ion and therefore assumes a spectrum intermediate between a red and a blue extreme. At the K stage the Schiff base nitrogen moves from  $\text{A}_3$  to  $\text{A}_1\text{H}$ , i.e., to the vicinity of an uncharged group. Hence, the spectrum is red shifted. At the L stage the group  $\text{A}_1$  deprotonates and becomes charged, a process which results in a blue shift of the spectrum. At the two M stages (only the second might be stable) the spectrum is necessarily moved to the extreme blue. At the N stage the chromophore interacts with a




FIGURE 5 Comparison of calculated and observed  $\text{C}_{14}\text{--}\text{C}_{15}$  stretch frequencies for the 13,14-*cis* model shown in Fig. 4. The experimental assignments for  $\text{K}_{625}$  are from Gerwert et al. (20), those for  $\text{O}_{640}$  are from Smith et al. (12) and for  $\text{M}_{412}$  it is our assignment according to a spectrum by Smith et al. (1). For the remaining assignments see the caption to Fig. 3.

charged group  $A_2^-$ , the interaction being very similar to that at the L stage. Hence, the N spectrum should be very similar to that of L. Since the rate constants of the transitions in this part of the cycle are all of similar magnitude, it might be difficult to separate N from L by time-resolved spectroscopy. The N intermediate re-isomerizes to the O stage (12) in which the situation arises that the Schiff base nitrogen sees an uncharged group  $A_3H$ . Hence, O should exhibit a red-shifted spectrum. The removal of a proton from  $A_3$  restores the initial  $Br_{568}$  stage.

The diagram in Fig. 4 also shows that side reactions shortcutting the cycle by means of slightly altered protonation-deprotonation reactions are possible. The shortcut would leave out the O intermediate. This implies the prediction that a strong correlation should exist between the appearance of O in the cycle and the yield of  $H^+$ 's pumped. Such correlation has, in fact, been observed by Renard et al. (35).

Fig. 4 attributes a distinct isomeric state to each stage of the pump cycle. The interactions of the chromophore with the indicated acids suggest that the retinal Schiff base should correspond to different categories in the nomenclature used above: we identify Br, L, and N with a 3-Å compound, K and O with a pure cation, and M with the unprotonated Schiff base. In Fig. 5 we compare for the assumed intermediates the calculated and the observed  $C_{14}-C_{15}$  stretch frequencies. Except for a systematic slight upshift of the calculated frequencies, Fig. 5 demonstrates a very clear parallelism between theory and observation. We take this parallelism as evidence that the geometries of retinal in the pump cycle have been assigned correctly. The small deviations remaining for the K and O intermediates can easily be rationalized in terms of a small perturbation of the cationic  $\pi$ -system by a local electric field near  $C_5$  (36).

We would like to note that the N intermediate may have been observed recently in RR experiments: Alshut and Stockburger observed at high pH a long-lived species with the same decay time as the  $M_{412}$  intermediate and with an RR spectrum very similar to that of L (personal communication). Assuming that this is a spectrum of N and that N is 13-*cis*, we have assigned in Fig. 3 A and B, the lowest two fingerprint vibrations to the  $C_{10}-C_{11}$  and  $C_{14}-C_{15}$  stretches. It would be worthwhile to identify the  $C_{14}-C_{15}$  vibration in this spectrum by repeating the experiment with  $^{13}C$  labeled retinal.

We wish to conclude that the stereodynamics of retinal in BR's pump cycle appears to be in agreement with our original suggestion (6). This suggestion has stated already that the stereodynamics determined here from the vibrational spectra imply that retinal plays the role of a proton switch in BR. However, the conclusion reached needs to be confirmed by further investigations. The most pressing questions in this respect are: (a) What is the energy of the  $C_{14}-C_{15}$  stretch vibration of  $L_{550}$ ? (b) Can other "fingerprint" vibrations elucidate structural details of the Schiff

base segment of retinal in BR? (c) Does an intermediate N exist and can its isomeric state be elucidated by RR or IR spectra? (d) Can the solid state NMR methods (32) resolve the structure of intermediates in the pump cycle?

We are grateful to K. Gerwert and F. Siebert for informing us about their experimental results prior to publication and for stimulating discussions. We would also like to thank D. Oesterhelt for encouragement and support. The contribution of H. Treutlein to the assignment of vibrational modes is gratefully acknowledged.

This project has been supported by the Deutsche Forschungsgemeinschaft (SFB-143 C1).

Received for publication 21 October 1985 and in final form 15 January 1985.

## REFERENCES

1. Smith, S. O., J. Lugtenburg, and R. Mathies. 1985. Determination of retinal chromophore structure in bacteriorhodopsin with resonance Raman spectroscopy. *J. Membr. Biol.* 85:95-109.
2. Smith, S. O., A. B. Myers, J. A. Pardo, C. Winkel, P. P. J. Mulder, J. Lugtenburg, and R. Mathies. 1984. Determination of retinal Schiff base configuration in bacteriorhodopsin. *Proc. Natl. Acad. Sci. USA.* 81:2055-2059.
3. Oesterhelt, D. 1976. Bacteriorhodopsin as an example of a light-driven proton pump. *Angew. Chem. Int. Ed. Engl.* 15:17-24.
4. Stoerkenius, W., and R. Bogomolni. 1982. Bacteriorhodopsin and the purple membrane of *Halobacteria*. *Annu. Rev. Biochem.* 51:587-616.
5. Stoerkenius, W., R. H. Lozier, and R. Bogomolni. 1979. Bacteriorhodopsin and related pigments of *Halobacteria*. *Biochim. Biophys. Acta.* 505:215.
6. Schulten, K., and P. Tavan. 1978. A mechanism for the light-driven proton pump of *Halobacterium halobium*. *Nature (Lond.)* 272:85-86.
7. Orlandi, G., and K. Schulten. 1979. Coupling of stereochemistry and proton donor-acceptor properties of a Schiff base. A model of a light-driven proton pump. *Chem. Phys. Lett.* 64:370-374.
8. Schulten, K., Z. Schulten, and P. Tavan. 1984. An isomerization model for the pump cycle of bacteriorhodopsin. In *Information and Energy Transduction in Biological Membranes*. L. Bolis, E. J. M. Helmreich, and H. Passow, editors. Alan R. Liss, Inc., New York. 113-131.
9. Tavan, P., K. Schulten, and D. Oesterhelt. 1985. The effect of protonation and electrical interactions on the stereochemistry of retinal Schiff bases. *Biophys. J.* 47:415-430.
10. Tavan, P., K. Schulten, W. Gärtner, and D. Oesterhelt. 1985. Substituents at the  $C_{13}$  position of retinal and their influence on the function of bacteriorhodopsin. *Biophys. J.* 47:349-355.
11. Smith, S. O., A. B. Myers, R. Mathies, J. A. Pardo, C. Winkel, E. M. M. van den Berg, and J. Lugtenburg. 1985. Vibrational analysis of the all-*trans* retinal protonated Schiff base. *Biophys. J.* 47:653-644.
12. Smith, S. O., J. A. Pardo, P. P. J. Mulder, B. Curry, J. Lugtenburg, and R. Mathies. 1983. Chromophore structure in bacteriorhodopsin's  $O_{640}$  photointermediate. *Biochemistry.* 22:6141-6148.
13. Oesterhelt, D., M. Meentzen, and L. Schuhmann. 1973. Reversible dissociation of the purple complex in bacteriorhodopsin and identification of 13-*cis* and all-*trans* retinal as its chromophores. *Eur. J. Biochem.* 40:453-463.
14. Lewis, A., J. Spoonhower, R. A. Bogomolni, R. H. Lozier, and W. Stoerkenius. 1974. Tunable laser resonance Raman spectroscopy of bacteriorhodopsin. *Proc. Natl. Acad. Sci. USA.* 71:4462-4466.



15. Marcus, M. A., and A. Lewis. 1977. Kinetic resonance Raman spectroscopy: dynamics of deprotonation of the Schiff base of bacteriorhodopsin. *Science (Wash. DC)*. 195:1328-1330.
16. Aton, B., A. G. Doukas, R. H. Callender, B. Becker, and T. G. Ebrey. 1977. Resonance Raman studies of the purple membrane. *Biochemistry*. 16:2995-2999.
17. Pettei, M. J., A. P. Judd, K. Nakanishi, R. Henselmann, and W. Stoeckenius. 1977. Identification of retinal isomers isolated from bacteriorhodopsin. *Biochemistry*. 16:1955-1959.
18. Schulten, K.. 1978. An isomerization model for the photocycle of bacteriorhodopsin. In *Energetics and Structure of Halophilic Microorganisms*. S. R. Caplan and M. Ginzburg, editors. Elsevier/North Holland Biomedical Press, Amsterdam. 331-334.
19. Nuss, M. C., W. Zinth, W. Kaiser, E. Kölling, and D. Oesterhelt. 1985. Femtosecond spectroscopy of the first events of the photochemical cycle in bacteriorhodopsin. *Chem. Phys. Lett.* 117:1-7.
20. Gerwert, K., F. Siebert, J. A. Pardoën, C. Winkel, and J. Lugtenburg. 1985. Vibrational analysis of chromophore-configurations in bacteriorhodopsin during the photo cycle using FTIR difference spectroscopy. In *Spectroscopy of Biological Molecules*. A. J. P. Alix, L. Bernard, and M. Manfait, editors. Wiley, Chichester. 373-375.
21. Curry, B. 1983. Ph.D. thesis. University of California, Berkeley, CA.
22. Curry, B., A. Broek, J. Lugtenburg, and R. Mathies. 1982. Vibrational analysis of all-*trans* retinal. *J. Am. Chem. Soc.* 104:5274-5286.
23. Dewar, J. S., and W. Thiel. 1977. Ground state of molecules. 38. The MNDO method. Approximations and parameters. *J. Am. Chem. Soc.* 99:4899-4906.
24. Tavan, P., and K. Schulten. 1984. The effect of protonation on the thermal isomerization of stilbazolium betaines. *Chem. Phys. Lett.* 110:191-195.
25. Steiner, U., M. H. Abd-El-Kader, P. Fischer, and H. E. A. Kramer. 1978. Photochemical *cis/trans* isomerization of a stilbazolium betaine. A protolytic/photochemical reaction cycle. *J. Am. Chem. Soc.* 100:3190-3197.
26. Panchenko, Yu. N., P. Pulay, and F. Török. 1976. Prediction of vibrational spectra by the CNDO/2 force method. II. The calculation of vibrational frequencies of *cis* and *trans* forms of glyoxal, acrolein and 1,3 butadiene. *J. Mol. Struct.* 34:283-289.
27. Schulten, K., U. Dinur, and B. Honig. 1980. The spectra of carbonium ions, cyanine dyes and protonated Schiff base polyenes. *J. Chem. Phys.* 73:3927-3935.
28. Honig, B., A. D. Greenberg, U. Dinur, and T. G. Ebrey. 1976. Visual-pigment spectra: implications of the protonation of the retinal Schiff base. *Biochemistry*. 15:4993-4999.
29. Derguini, F., C. G. Caldwell, M. G. Motto, V. Balogh-Nair, and K. Nakanishi. 1983. Bacteriorhodopsins containing cyanine dye chromophores. Support for the external point charge model. *J. Am. Chem. Soc.* 105:646-648.
30. Nakanishi, K., V. Balogh-Nair, M. Arnaboldi, K. Tsujimoto, and B. Honig. 1980. An external point-charge model for bacteriorhodopsin to account for its purple color. *J. Am. Chem. Soc.* 102:7945-7947.
31. Kakitani, T., H. Kakitani, B. Honig, and K. Nakanishi. 1983. Symmetric charge distribution in the bacteriorhodopsin binding site. *J. Am. Chem. Soc.* 105:648-650.
32. Harbison, G. S., S. O. Smith, J. A. Pardoën, C. Winkel, J. A. Lugtenburg, J. A. Herzfeld, R. Mathies, and R. G. Griffin. 1984. Dark-adapted bacteriorhodopsin contains 13-*cis*,15-*syn* and all-*trans*,15-*anti* retinal Schiff bases. *Proc. Natl. Acad. Sci. USA*. 81:1706-1709.
33. Braiman, M., and R. Mathies. 1982. Resonance Raman spectra of bacteriorhodopsin primary photoproduct: evidence for a distorted 13-*cis* retinal chromophore. *Proc. Natl. Acad. Sci. USA*. 79:403-407.
34. Engelhard, M., K. Gerwert, B. Hess, W. Kreutz, and F. Siebert. 1985. Light-driven protonation changes of internal aspartic acids of bacteriorhodopsin: an investigation by static and time-resolved infrared difference spectroscopy using [4-<sup>13</sup>C]aspartic acid labeled purple membrane. *Biochemistry*. 24:400-407.
35. Renard M., P. Thirion, and M. Delmelle. 1983. Photoacoustic spectroscopy of bacteriorhodopsin photocycle. *Biophys. J.* 44:211-218.
36. Spudich, J. L., D. A. McCain, K. Nakanishi, M. Okabe, N. Shimizu, H. Rodman, B. Honig, and R. A. Bogomolni. 1986. Chromophore/protein interaction in bacterial sensory rhodopsin and bacteriorhodopsin. *Biophys. J.* 49:479-483.