

Markus Dittrich and Klaus Schulten. Zooming in on ATP hydrolysis in F_1 . J. Bioenerg. Biomembr., 37:441-444, 2005.

Zooming in on ATP hydrolysis in F_1

Markus Dittrich¹ and Klaus Schulten¹

¹*Theoretical and Computational Biophysics Group, Beckman Institute, University of Illinois at Urbana-Champaign, 405 N. Mathews, Urbana, IL 61801, USA*

Corresponding author: Klaus Schulten,
Beckman Institute, University of Illinois at Urbana-Champaign,
405 N. Mathews Ave., Urbana, IL 61801
Tel: (217) 244-1604 Fax: (217) 244-6078
E-mail: kschulte@ks.uiuc.edu

Abstract

We summarize our current view of the reaction mechanism in F_1 -ATPase as it has emerged from experiment, theory, and computational studies over the last several years. ATP catalysis in the catalytic binding pockets of F_1 takes place without the release of any significant free energy and is efficiently driven by the combined action of two water molecules utilizing a so-called protein relay-mechanism. The chemical reaction itself is controlled by the spatial position of a key arginine residue.

Key words: F_1 -ATPase; ATP; hydrolysis; QM/MM; molecular mechanics

1 Introduction

In the decade that has passed since the first X-ray crystal structure of F_1 was determined in 1994 by Abrahams et al. (1994), our understanding of the microscopic processes underlying F_1 function has advanced steadily. In addition to the vast amount of information contained in the structures themselves, this advance can be attributed also to the fact that biochemical experiments may now be interpreted in a structural context. The atomic resolution data permit computer simulations to study F_1 function, e.g., stalk rotation (Böckmann and Grubmüller, 2002) or the ATP hydrolysis reaction in the catalytic sites (Dittrich et al., 2003; Strajbl et al., 2003; Dittrich et al., 2004).

Our present understanding of the chemo-mechanical coupling in F_1 -ATPase, i.e., the coupling of ATP hydrolysis to stalk rotation during F_1 's rotatory catalytic cycle, is still incomplete. However, recent experiments (Nishizaka et al., 2004; Shimabukuro et al., 2003) and computer simulations (Dittrich et al., 2003; Strajbl et al., 2003; Dittrich et al., 2004) have provided a much clearer picture of the underlying hydrolysis events, particularly their energetics and their coupling to the protein environment. It can be hoped, therefore, that further investigation will lead to a more complete understanding of how F_1 achieves its coupling of ATP synthesis/hydrolysis to stalk rotation and proton current through F_o .

Here, we will summarize our understanding of the ATP hydrolysis reaction in the catalytic sites of F_1 -ATPase and its coupling to protein conformational changes that follows from recent computer simulations as well as experimental findings and theoretical modeling. To assist our discussion, we will refer to "ATP hydrolysis" as the overall reaction process starting from an ATP molecule in solution and ending with ADP and P_i in solution. We will use the term "ATP catalysis" to describe the actual chemical bond breaking event in the catalytic sites of F_1 -ATPase.

2 The role of ATP catalysis

The ATP hydrolysis reaction powering F_1 and many other molecular motors and cellular processes is exothermic with a ΔG_0 of approximately -7kcal/mol (Berg et al., 2002). As single-molecule experiments (Nishizaka et al., 2004; Shimabukuro et al., 2003) have shown, the ATP hydrolysis reaction driving the rotatory catalytic cycle of F_1 can be divided into at

least three distinct events: ATP binding, ATP catalysis, and unbinding of products ADP and P_i . In principle, each of these processes individually or any combination of them could be the force generating step, i.e., the one in which the free energy stored in ATP is being extracted by the protein and converted into mechanical torque acting on the F_1 stalk. Experimentally, both ATP catalysis (Weber et al., 2000) and the initial binding of ATP to the catalytic sites (Nishizaka et al., 2004; Shimabukuro et al., 2003) have been suggested as force generating steps.

From an energetic point of view, however, it seems unlikely that the protein is able to extract, in an efficient manner, the free energy released during ATP catalysis, since the latter is an ultra-fast process taking place on a fs time scale. This was recently corroborated by computer simulations (Dittrich et al., 2003, 2004) which revealed that no net free energy is released during ATP catalysis itself. In their paper (Dittrich et al., 2003, 2004), the authors showed that the catalysis reaction energy profile changed from strongly endothermic in the β_{TP} catalytic site to approximately equi-energetic in β_{DP} . This leaves either reactant binding (ATP), product unbinding (ADP, P_i), or both as possible candidates for force generation.

Considering the charged nature of the Mg-ATP complex and the multitude of possible hydrogen bond/salt bridge interactions of ATP's tri-phosphate moiety with the protein, an ATP binding-induced forced closure of the previously open β_E pocket as the force generating step seems very attractive. This type of mechanism has been proposed by Oster and coworkers (Oster and Wang, 2000) in their theoretical modeling studies and termed *ATP binding zipper*.

Assuming that the tight binding of ATP into the binding pockets is the actual force generating event during rotatory catalysis and taking into account the fact that no free energy is released during ATP catalysis, it follows that the unbinding of products ADP and P_i from β_{DP} will actually require the input of free energy. This energy could, e.g., be supplied by ATP binding to the neighboring empty β_E site. Oster et al. have suggested an elastic recoil driven by energy stored during the initial ATP binding event as free energy source for P_i unbinding (Oster and Wang, 2000). However, it is not clear how proteins could achieve such a localized and efficient storage of elastic energy without much dissipation.

What, then, is the role of the actual chemical step if not force generation? Computational mutation studies (Dittrich et al., 2004) have shown that the spatial position of important residues in the catalytic binding sites, in particular the arginine finger residue $\alpha R373$, can switch on and off catalysis by means of adjusting the reaction energy barrier and profile. More importantly, this switch mechanism can enforce cooperativity between the catalytic sites. For example, F_1 might inhibit ATP hydrolysis in β_{TP} via retracting the guanidinium group of $\alpha R373$ from the bound nucleotide until solution ATP binds to the empty β_E site. This binding event causes movement of $\alpha R373$ back into the β_{TP} catalytic site, thereby, re-enabling ATP hydrolysis and the progression of the rotatory catalytic cycle, effectively establishing a line of communication between the β_E and β_{TP} catalytic sites.

3 Efficient ATP hydrolysis in F_1 involves two water molecules

ATP hydrolysis in an aqueous environment is initiated by nucleophilic attack of a single water molecule (or hydroxide anion) on the γ -phosphate group of the nucleotide. A similar mechanism was also assumed to hold inside the catalytic binding sites of F_1 -ATPase and other ATP hydrolyzing enzymes. Support for this view came from the presence of electron density for a single water molecule in the first X-ray crystal structure of F_1 (Abrahams et al., 1994), that was located between the γ -phosphate group and the carboxyl group of $\beta E188$. The latter was assumed to polarize the nucleophilic water molecule or to act as a general base, thereby, facilitating nucleophilic attack (Abrahams et al., 1994).

In contrast, recent computational studies employing combined quantum mechanical/molecular mechanical simulations (Dittrich et al., 2003, 2004) revealed that ATP catalysis and nucleophilic attack in the catalytic binding sites of F_1 involves two water molecules working together in a concerted fashion. This is illustrated in Figure 1 which shows the reactant, transition, and product state conformations during the ATP hydrolysis reaction in the β_{DP} binding site of F_1 . Figure 2 shows a schematic view of this so-called multi-center *protein-relay mechanism*, in which a second water molecule, WAT2, abstracts a proton from the nucleophilic water, WAT1, thereby, acting as a base. This leads to an excess electron density on WAT1-O, facilitating nucleophilic attack on ATP- P_γ . The abstracted proton on WAT2 contributes to the formation of an hydronium ion-like construct in the transition state and is electrostatically stabilized by the protein environment, predominantly the negatively charged carboxyl group of $\beta E188$. This is shown schematically in Figure 2.

The electron density transferred from the nucleophilic WAT1 to the terminal phosphate group upon bond formation in turn facilitates protonation of ATP- O_γ by the hydronium ion. Interestingly, this protonation event does not involve the proton originally abstracted from WAT1, but rather one that stems from WAT2.

Dittrich et al. (2004) have shown that such a protein-relay mechanism is energetically favorable compared to nucleophilic attack of a single water molecule. This is mainly due to the fact that the involvement of a second water molecule offers a very efficient mechanism for nucleophile polarization and proton transfer from the nucleophile to ATP. As shown in Dittrich et al. (2004), the proton relay mechanism allows the nucleophilic WAT1 to perform an in-line attack on P_γ giving rise to a WAT1-O – ATP- P_γ separation in the transition state that is only slightly longer (0.2Å) than the final bonding distance without any distortion of the planar penta-covalent arrangement of the terminal phosphate group. This facilitates the efficient transfer of electron density from the nucleophile to the site of protonation on one of the γ -phosphate oxygens and, consequently, assists proton transfer. In contrast, the nucleophilic attack of a single water molecule and the associated direct proton transfer to the γ -phosphate group requires the system to assume a non-inline and distorted penta-covalent transition state conformation. This leads to a significantly longer WAT1-O – ATP- P_γ separation compared to the final bonding distance ($\sim 0.5\text{\AA}$), making proton transfer via the latter mechanism less efficient and energetically unfavorable.




Figure 1: ATP hydrolysis reaction in the β_{DP} catalytic site. Shown are the reactant (**a**), transition (**b**), and final (**c**) state conformations. In (**b**) the involvement of two water molecules is clearly discernible in the form of an intermediate H_3O^+ -like construct. (Figure adapted with permission from Dittrich et al. (2004). Copyright 2004 Biophysical Society.)

In this context it is important to realize that such a proton-relay mechanism is made possible exclusively by the protein environment provided by the binding pocket and does




Figure 2: Schematic representation of the multi-center reaction for ATP hydrolysis, showing the a) reactant, b) transition, c) intermediate and d) product state. (Figure reprinted with permission from Dittrich et al. (2003). Copyright 2003 Biophysical Society.)

not exist in solution, making this a genuine enzymatic pathway.

Several other studies have found evidence for the involvement of such proton relays for efficient nucleotide hydrolysis. Scheidig et al. (1999) proposed the participation of two water molecules during GTP hydrolysis in p21^{ras} on the basis of X-ray crystal structure data. A recent theoretical study by Li and Cui (2004) investigating ATP hydrolysis in myosin also identified a proton-relay mechanism involving the hydroxy group of a serine residue as the energetically most favorable one. Finally, an involvement of two water molecules during ATP catalysis in F₁ is also consistent with recent experimental mutation studies (Ahmad and Senior, 2004), which found that residue βN243 in F₁ is involved in the orientation of the attacking and a second, associated water molecule, which could correspond to WAT1 and WAT2 in the computational studies of Dittrich et al. (2004).

4 Conclusions

Over the last years, the combination of experiment, theory, and computer simulation have provided new insights into the microscopic events during the chemo-mechanical coupling in F₁-ATPase. It has emerged that the actual ATP catalysis reaction does not release any significant free energy to drive stalk rotation, but rather furnishes a control mechanism governed by the spatial position of the arginine finger residue αR373 enabling catalytic site cooperativity. Efficient ATP hydrolysis is catalyzed by the cooperative action of two water molecules giving rise to rapid nucleophilic attack and proton transfer from the nucleophilic water to ATP. As studies of different ATP hydrolyzing enzymes have shown, this two-water

mechanism can be expected to be of fundamental importance for protein catalyzed nucleotide hydrolysis in systems other than F_1 .

5 Acknowledgments

This work is supported by grants from the National Institutes of Health PHS-5-P41-RR05969 and the National Science Foundation MCB02-34938. The authors gladly acknowledge supercomputer time provided by Pittsburgh Supercomputer Center and the National Center for Supercomputing Applications via National Resources Allocation Committee grant MCA93S028.

References

- Abrahams, J., A. Leslie, R. Lutter, and J. Walker. 1994. Structure at 2.8 Å resolution of F_1 -ATPase from bovine heart mitochondria. *Nature* 370:621–628.
- Ahmad, Z. and A. E. Senior. 2004. Role of β Asn-243 in the phosphate-binding subdomain of catalytic sites of *Escherichia coli* F_1 -ATPase. *J. Biol. Chem.* 44:46057–46064.
- Berg, J. M., J. L. Tymoczko, and L. Stryer. 2002. Biochemistry. W. H. Freeman and Co., New York, fifth edition.
- Böckmann, R. A. and H. Grubmüller. 2002. Nanoseconds molecular dynamics simulation of primary mechanical energy transfer steps in F_1 -ATP synthase. *Nature Struct. Biol.* 9:198–202.
- Dittrich, M., S. Hayashi, and K. Schulten. 2003. On the mechanism of ATP hydrolysis in F_1 -ATPase. *Biophys. J.* 85:2253–2266.
- Dittrich, M., S. Hayashi, and K. Schulten. 2004. ATP hydrolysis in the β_{TP} and β_{DP} catalytic sites of F_1 -ATPase. *Biophys. J.* 87:2954–2967.
- Li, G. and Q. Cui. 2004. Mechanochemical coupling in myosin: A theoretical analysis with molecular dynamics and combined QM/MM reaction path calculations. *J. Phys. Chem. B* 108:3342–3357.
- Nishizaka, T., K. Oiwa, H. Noji, S. Kimura, E. Muneyuki, M. Yoshida, and K. Kinoshita Jr. 2004. Chemomechanical coupling in F_1 -ATPase revealed by simultaneous observation of nucleotide kinetics and rotation. *Nat. Struct. Mol. Biol.* 11:142–148.
- Oster, G. and H. Wang. 2000. Reverse engineering a protein: the mechanochemistry of ATP synthase. *Biochim. Biophys. Acta* 1458:482–510.

- Scheidig, A. J., C. Burmester, and R. S. Goody. 1999. The pre-hydrolysis state of p21^{ras} in complex with GTP: new insights into the role of water molecules in the GTP hydrolysis reaction of ras-like proteins. *Structure* 7:1311–1324.
- Shimabukuro, K., R. Yasuda, E. Muneyuki, K. Y. Hara, K. Kinosita Jr., and M. Yoshida. 2003. Catalysis and rotation of F₁ motor: Cleavage of ATP at the catalytic site occurs in 1 ms before 40° substep rotation. *Proc. Natl. Acad. Sci. USA* 100:14731–14736.
- Strajbl, M., A. Shurki, and A. Warshel. 2003. Converting conformational changes to electrostatic energy in molecular motors: The energetics of ATP synthase. *Proc. Natl. Acad. Sci. USA* 100:14834–14839.
- Weber, J., S. Nadanaciva, and A. E. Senior. 2000. ATP-driven rotation of the γ subunit in F₁-ATPase. *FEBS Lett.* 483:1–5.